



# Novel sodium channel gene mutations in *Blattella germanica* reduce the sensitivity of expressed channels to deltamethrin

J. Tan<sup>1,a</sup>, Z. Liu<sup>1 a</sup>, T.-D. Tsai<sup>b</sup>, S.M. Valles<sup>c</sup>, A.L. Goldin<sup>d</sup>, K. Dong<sup>a,\*</sup>

<sup>a</sup> Department of Entomology and Neuroscience Program, Michigan State University, East Lansing, MI 48824, USA

<sup>b</sup> Pharmacology, Pharmacia Corp., Kalamazoo, MI 49007, USA

<sup>c</sup> USDA-ARS, Center for Medical, Agricultural and Veterinary Entomology, Gainesville, FL 32608, USA

<sup>d</sup> Department of Microbiology and Molecular Genetics, University of California, Irvine, CA 92697, USA

Received 18 January 2001; received in revised form 28 May 2001; accepted 10 July 2001

---

## Abstract

Pyrethroid insecticides alter the normal gating of voltage-gated sodium channels in the nervous system. Three sodium channel mutations (E434K, C764R, L993F) were recently identified in pyrethroid resistant German cockroach populations. In this report, we show that the L993F mutation decreased sodium channel sensitivity to the pyrethroid, deltamethrin, by five-fold in *Xenopus* oocytes. In contrast, neither E434K nor C764R alone decreased channel sensitivity to deltamethrin. However, E434K or C764R combined with L993F reduced deltamethrin sensitivity by 100-fold. Furthermore, concomitant presence of all three mutations (KRF) reduced channel sensitivity to deltamethrin by 500-fold. None of the mutations significantly affected channel gating. However, sodium current amplitudes from the mutant sodium channel carrying either E434K or C764R alone were much reduced compared to those of the wild-type channel or the channel carrying the double or triple mutations (KF, RF and KRF). These results indicated that evolution of sodium channel insensitivity in the German cockroach is achieved by sequential selection of a primary mutation L993F and two secondary mutations E434K and C764R, and concomitant presence of all three mutations dramatically reduced sodium channel sensitivity to deltamethrin. © 2002 Elsevier Science Ltd. All rights reserved.

**Keywords:** Knockdown resistance; Pyrethroids; Insecticide resistance; Sodium channel; *Xenopus* oocyte expression system

---

## 1. Introduction

The voltage-dependent sodium channel is a major target for a variety of neurotoxins including DDT and pyrethroid insecticides (Narahashi, 1986; Catterall, 1988; Hille, 1992; Zlotkin, 1999). These neurotoxins have well-defined effects on many functional properties of the sodium channel, such as conductance, ion selectivity, activation, and inactivation (Catterall, 1988). They are widely used as molecular probes for characterization of functional regions within sodium channel subunits

(Catterall, 1992). Indeed, these neurotoxins are often classified according to their unique binding sites on the sodium channel (Catterall, 1988). For example, both DDT and pyrethroid insecticides, although structurally different, cause repetitive discharge or nerve membrane depolarization by modifying sodium channel gating (references in Narahashi, 1988; Soderlund and Bloomquist, 1989). Results from electrophysiological (such as Narahashi, 1988; Takeda and Narahashi, 1988; Song et al., 1996) and pharmacological studies (such as Jacques et al., 1980, Bloomquist and Soderlund, 1988; Brown et al., 1988; Lombet et al., 1988; Trainer et al., 1997) suggest a distinct receptor site(s) for pyrethroids. Although the electrophysiological effects of pyrethroids on sodium channels have been well-defined, our understanding of the molecular interactions between sodium channels and pyrethroids remains rudimentary. Because pyrethroids are highly lipophilic, pharmacological studies have been either impossible to perform or of limited utility due to very high levels of nonspecific ligand binding. Identifi-

---

**Abbreviations:** DDT, 1,1,1-trichloro-2,2-bis(*p*-chlorophenyl)ethane; kdr, knockdown resistance; TTX, tetrodotoxin.

\* Corresponding author. Rm. 106, Center for Integrated Plant Systems, Michigan State University, East Lansing, MI 48824, USA. Tel.: +1-517-432-2034; fax: +1-517-353-5598.

**E-mail addresses:** tanj@msu.edu (J. Tan), dongk@pilot.msu.edu (K. Dong).

<sup>1</sup> These authors contributed equally to this work.

cation and electrophysiological characterization of mutations that affect sodium channel sensitivity to pyrethroids provide a powerful complementary approach, and should lead to a better understanding of sodium channel function and of the molecular interactions between sodium channels and pyrethroids that act on them.

Important sources of mutations that affect sodium channel interactions with DDT and pyrethroids are DDT- and pyrethroid-resistant insects. The intense use of these compounds has led to the selection of resistant populations in many insects. One important resistance mechanism is the called knockdown resistance (kdr) in housefly and kdr-type resistance in German cockroaches. It confers resistance to both knockdown (rapid paralysis) and killing by pyrethroids through reduced neuronal sensitivity to these compounds (references in Soderlund and Bloomquist, 1990). The kdr or kdr-type phenotype is associated with a leucine to phenylalanine mutation (L1014F in houseflies, L993F in cockroaches) in IIS6 of the insect Para-homologous sodium channel protein in about a dozen insect species (Williamson et al., 1996; Miyazaki et al., 1996; Dong, 1997; Martinez-Torres et al. 1997, 1998; Guerrero et al., 1997; Martinez-Torres et al., 1999a,b; Lee et al., 1999a). The super-kdr resistance phenotype in the house fly is associated with both the L1014F mutation in IIS6 and an M to T mutation (M918T) in the linker region between S4 and S5 of domain II (Williamson et al., 1996). In pyrethroid-resistant tobacco budworm (*Heliothis virescens*), the L1029H mutation in IIS6 (corresponding to L993F in cockroaches) or a V421M mutation in IS6 is associated with pyrethroid resistance (Park and Taylor, 1997; Park et al., 1997; Lee et al., 1999b).

Using the *Xenopus* oocyte expression system, Smith et al. (1997) showed that the house fly sodium channel carrying the L1014F mutation alone was 10-fold less sensitive to the pyrethroid cismethrin than the wild-type channel. The super-kdr M918T mutation, when combined with the L1014F mutation, further reduced sodium channel sensitivity to cismethrin (Lee et al., 1999c). Introduction of the two house fly mutations into the *Drosophila* Para protein greatly decreased channel sensitivity to deltamethrin (Vais et al., 2000). The V421M and L1029H sodium channel mutations associated with pyrethroid resistance in *Heliothis virescens* were also confirmed to reduce sodium channel sensitivity to the pyrethroid permethrin (Zhao et al., 2000). These studies clearly established that these sodium channel mutations play a significant role in sodium channel insensitivity to pyrethroids.

The L993F mutation was also identified in more than twenty pyrethroid-resistant German cockroach populations (Miyazaki et al., 1996; Dong, 1997; Dong et al., 1998). However, the house fly super-kdr mutation was not found in any of the five highly resistant German

cockroach populations (Dong et al., 1998). Instead, four novel *para* mutations (D58G, E434K, C764R and P1880L) were found to be associated with high-level pyrethroid resistance in this insect species (Liu et al., 2000). This raises the intriguing possibility that different *para*<sup>CSMA</sup> gene mutations may have been selected in different insect species for high levels of pyrethroid resistance. In this study, we investigate the effects of E434K, C764R, and L993F mutations, which were present in all five highly resistant strains described by Liu et al. (2000), on cockroach Para<sup>CSMA</sup> channel properties and sensitivity to a pyrethroid insecticide, deltamethrin.

## 2. Materials and methods

### 2.1. Polymerase chain reaction (PCR) amplification and cloning of a full-length *para* cDNA and *D. melanogaster tipE* cDNA

The nucleotide sequence of the coding region of the cockroach sodium channel gene *para*<sup>CSMA</sup> was determined previously by sequencing overlapping cDNA clones (Dong, 1997). In this study, mRNA was isolated from the heads and thoraces of adult male German cockroaches of strain CSMA, which is susceptible to pyrethroids and DDT. The methods for RNA isolation and cDNA synthesis are the same as described in Dong (1997). For amplification of *para*<sup>CSMA</sup> cDNA, the first strand cDNA was synthesized using a primer corresponding to the sequence in the 3' untranslated region (5'-AAGAATGAGGCATCTCCGAG-3'). The entire coding region (ca. 6 kb) of *para*<sup>CSMA</sup> was amplified by PCR using the eLONGase enzyme mix (Gibco/BRL, Rockville, MD). The amplification reaction mixture (50  $\mu$ l) contained 0.5  $\mu$ l cDNA, 0.2  $\mu$ mol 5' primer (5'-CCGGTACCGCCACCATGGCCGACGACTCGTCCTCAATCTCAGA-3'; Kozak sequence italics), 0.2  $\mu$ mol 3' primer (5' -CCGGATCCAATCAAGCGAAGATGTGAGA-3'), 200  $\mu$ M each dNTP, 1 U of eLONGase, 1.5 mM MgCl<sub>2</sub>, and 1 $\times$  PCR reaction buffer supplied by the manufacturer. To facilitate cloning, *Kpn*I and *Bam*H1 restriction site sequences were added to the 5' and 3' primers, respectively. The sequence of the 5' primer was designed to conform to the Kozak sequence for high-efficiency translation. The T<sup>+4</sup> to G<sup>+4</sup> change results in the Ser to Ala substitution. The PCR amplification was performed for 30 cycles of 30 s at 94°C, 30 s at 58°C, and 7 min at 68°C followed by incubation at 68°C for 10 min. The amplified full-length cDNA with *Kpn*I and *Bam*H1 sites attached to the 5' and 3' ends, respectively, was cloned into pGH19 (kindly provided by Dr B. Ganetzky, University of Wisconsin, Madison). The entire insert of one clone, KD1, was sequenced. The *D. melanogaster tipE* cDNA was amplified and cloned into

pBluescript KS(+) using PCR primers based on the published sequence (Feng et al., 1995).

## 2.2. Site-directed mutagenesis

KD1 was mutagenized to generate a series of mutant constructs with various combinations of the three mutations, E434K, C764R, and L993F. Specifically, a 2.3-kb *NdeI* fragment containing all three nucleotides to be mutated was excised from KD1 and cloned into pAlter 1 (Promega Corp., Madison, WI) for site-directed mutagenesis. Point mutations were introduced using the Altered Sites II in vitro Mutagenesis System (Promega Corp.) and confirmed by DNA sequencing. The mutated *NdeI* fragments were excised from pAlter1 and cloned back into pGH19 containing the remainder of the KD1 sequence to produce the corresponding mutant constructs.

## 2.3. Expression of *Para*<sup>CSMA</sup> sodium channels in *Xenopus* oocytes

Oocytes were obtained surgically from oocyte-positive female *Xenopus laevis* (Nasco, Ft. Atkinson, WI) and incubated with 1 mg/ml Type IA collagenase (Sigma Co., St. Louis, MO) in Ca<sup>2+</sup>-free ND 96 medium, which contains 96 mM NaCl, 2 mM KCl, 1 mM MgCl<sub>2</sub>, and 5 mM HEPES, pH 7.5. Follicle cell remaining on the oocytes were removed with forceps. Isolated oocytes were incubated in ND-96 medium containing 1.8 mM CaCl<sub>2</sub> supplemented with 50 µg/ml gentamicin, 5 mM pyruvate, and 0.5 mM theophylline (Goldin, 1992). Healthy stage V–VI oocytes were used for cRNA injection. To prepare cRNA for oocyte injection, plasmid DNA of the *para* or *tipE* construct was linearized with *NotI*, which does not cut the insert, followed by in vitro transcription with T7 polymerase using the mMACHINE kit (Ambion). *para*<sup>CSMA</sup> cRNA (1 ng) and *tipE* cRNA (1 ng) were co-injected into oocytes, which were incubated at 19°C for three–seven days.

## 2.4. Electrophysiological recording and analysis

Methods for electrophysiological recording and data analysis are similar to those described previously (Kontis and Goldin, 1993). Sodium currents were recorded using standard two-electrode voltage clamping. The borosilicate glass electrodes were filled with filtered 3 M KCl and had resistance less than 0.5 MΩ. Currents were measured using the oocyte clamp instrument OC725C (Warner Instrument Corp. Hamden, CT), Digidata 1200A interface (Axon Instrument, Foster City, CA) and pCLAMP 6 software (Axon Instrument Foster City, CA). All experiments were performed at room temperature (20–22°C). Capacitive transient and linear leak currents were corrected using P/N subtraction technique or

by subtraction of records obtained in the presence of 20 nM tetrodotoxin (TTX), which completely blocks Para sodium channels.

For application of deltamethrin, the disposable perfusion system developed by Tatebayashi and Narahashi (1994) was used. Briefly, the test solution was transferred into a Petri dish placed on a supportstand. Two glass capillary tubes (10 cm in length) connected together with a short length of Tygan tubing were used to aid solution flow from the Petri dish to the recording chamber. The solution flow was controlled by hydrostatic force created by adjusting the level of the Petri dish relative to the recording chamber. Disposable recording chambers (1–1.5 ml volume) were made with glue dams on glass slides. Because deltamethrin is extremely lipophilic, recording chambers, perfusion system, and the glass agarose bridges connecting the oocyte chamber with the ground electrode chamber were all discarded after a single use. The deltamethrin stock solution (100 mM) was prepared in dimethylsulfoxide (DMSO). The working solutions were made in ND-96 medium immediately before use. Effects of deltamethrin on sodium channel tail currents reached a steady-state level within 5 min after perfusion.

The voltage-dependence of sodium channel conductance (*G*) was calculated by measuring the peak current at test potentials ranging from –120 to +60 mV in 5 mV increments and dividing by (*V* – *V*<sub>rev</sub>), where *V* is the test potential and *V*<sub>rev</sub> is the reversal potential for sodium. Reversal potentials were determined from the *I*–*V* curves. Peak conductance values were fit with a two-state Boltzmann equation of the form  $G = 1 - [1 + \exp((V - V_{1/2})/k)]^{-1}$ , in which *V* is the potential of the voltage pulse, *V*<sub>1/2</sub> is the half-maximal voltage for activation, and *k* is the slope factor.

The voltage-dependence of sodium channel inactivation was determined using 200 ms inactivating pre-pulses from a holding potential of –120 to +40 mV in 5 mV increments, followed by test pulses to –5 mV for 12 ms. The peak current amplitude during the test depolarization was normalized to the maximum current amplitude, and plotted as a function of the pre-pulse potential. The data were fit with a two-state Boltzmann equation of the form  $I = I_{\max} * [1 + (\exp(V - V_{1/2})/k)]^{-1}$ , where *I*<sub>max</sub> is the maximal current evoked, *V* is the potential of the voltage pulse, *V*<sub>1/2</sub> is the voltage at which 50% of the current is inactivated (the midpoint of the inactivation curve), and *k* is the slope factor.

The percentage of channels modified by deltamethrin was calculated using the equation  $M = [I_{\text{tail}} / (E_h - E_{\text{Na}})] / [I_{\text{Na}} / (E_t - E_{\text{Na}})] \times 100$  (Tatebayashi and Narahashi, 1994), in which *I*<sub>tail</sub> is the maximal tail current amplitude, *E*<sub>h</sub> is the potential to which the membrane is repolarized, *E*<sub>Na</sub> is the reversal potential for sodium current determined from the *I*–*V* curve, *I*<sub>Na</sub> is the amplitude of

the peak current during depolarization before deltamethrin exposure, and  $E_t$  is the potential of step depolarization. The concentration–response data were fitted to the Hill equation:  $M = M_{\max} / (1 + (EC_{50} / [\text{deltamethrin}])^n)$ , where [deltamethrin] and  $EC_{50}$  represent the concentration of deltamethrin and the concentration to produce the half-maximal effect, respectively,  $n$  represents the Hill coefficient, and the  $M_{\max}$  is the maximal percentage of sodium channels modified.

### 3. Results

#### 3.1. Functional expression of *Para*<sup>CSMA</sup> in *Xenopus* oocytes

We isolated a full-length cockroach *para*<sup>CSMA</sup> cDNA clone, KD1, by RT-PCR. Four amino acid sequence discrepancies, R502G, L1285P, V1685A and I1806L were found between KD1 and the previously published sequence (Dong, 1997). KD1 contains alternative exon 1 but not alternative exon 2 and another alternative exon (<sup>727</sup>VSIYYFPT<sup>735</sup>, Dong, 1997). Expression of KD1 in *Xenopus* oocytes confirmed that KD1 encodes a functional sodium channel protein. Specifically, sodium currents with a peak current of approximately 2–5  $\mu$ A were detected on day 4 or 5 in most oocytes co-injected with *para*<sup>CSMA</sup> and *tipE* cRNA (1 ng each). The sodium currents were completely inhibited by 10 nM TTX (Fig. 1B). Sodium current traces from a representative oocyte are shown in Fig. 1A. The oocyte was clamped at a holding potential of –120 mV and depolarized to test potentials between –80 and 60 mV in 5 mV increments. Sodium current through the cockroach *Para*<sup>CSMA</sup> channel was detected at approximately –40 mV and reached peak values at –10 mV (Fig. 1C). The average relative sodium conductance was calculated as described in Section 2 and plotted as a function of depolarizing test potentials (Fig. 1D). The voltage for half-maximal activation was  $-29.0 \pm 1.3$  mV and the slope factor was  $5.5 \pm 0.5$  (Table 1). To determine the voltage-dependence of steady-state inactivation, oocytes were held at –120 mV and depolarized with a series of 200-ms inactivating prepulses from –120 to +40 mV in 5 mV increments, followed by 12-ms test pulses to –5 mV to measure channel availability. The voltage-dependence of steady-state inactivation was obtained by plotting the normalized peak current as a function of prepulse potentials, as described in Section 2 (Fig. 1D). The voltage for half-maximal inactivation was  $-42.4 \pm 0.4$  mV and the slope factor was  $5.0 \pm 0.1$  (Table 1).

#### 3.2. Effects of *kdr*-type mutations on channel expression and gating properties

Five *para*<sup>CSMA</sup> mutations (D58G, E434K, C764R, L993F, and P1880L) were found to be associated with

*kdr*-type resistance in the German cockroach (Liu et al., 2000). The L993F mutation was detected in twenty of twenty four recently field-collected pyrethroid-resistant populations (Dong, 1997; Dong et al., 1998). The E434K and C764R mutations were found to coexist with the L993F mutation in only the five most highly pyrethroid-resistant populations (Liu et al., 2000). The D58G and P1880L mutations were found in only one of the five populations (Liu et al., 2000). We hypothesized that the L993F mutation is responsible for a low level of knock-down resistance in cockroaches, whereas the other four mutations are responsible for high levels of *kdr*-type resistance. To test this hypothesis, we generated constructs with all eight possible combinations of the three most common mutations, E434K, C764R and L993F (Fig. 2). For simplicity, the mutants will be referred to by the single letter code for the new amino acid that is present, with multiple letters indicating the presence of multiple mutations, as listed in Table 1. We first examined the effects of the various combinations of mutations on the peak current of the sodium channel in oocytes. Mutant channels, *Para*<sup>CSMA-F</sup>, *Para*<sup>CSMA-KR</sup>, *Para*<sup>CSMA-RF</sup>, *Para*<sup>CSMA-KF</sup> and *Para*<sup>CSMA-KRF</sup> all expressed sodium currents with peak currents comparable to that of the wild-type channel (Table 2). The peak currents of mutant channels, *Para*<sup>CSMA-K</sup> and *Para*<sup>CSMA-R</sup>, however, were significantly (four- to five-fold) lower than that of the wild-type channel (Table 2).

Effects of mutations on voltage-dependence of activation and steady-state inactivation were also examined. Sodium currents recorded in oocytes expressing mutant channels exhibited voltage-dependence of activation and inactivation similar to those of the wild-type channel (Table 1), demonstrating that the gating properties of the mutant channels are apparently normal.

#### 3.3. *kdr*-type Mutations altered the *Para*<sup>CSMA</sup> channel sensitivity to deltamethrin

Pyrethroids alter sodium channel function by slowing inactivation and deactivation (Narahashi, 1988). Specifically, the sodium current during depolarization is prolonged, and the tail current associated with repolarization is greatly augmented. The pyrethroid-induced tail current has been used to quantify pyrethroid modification of the sodium channel (Lund and Narahashi, 1982; Vijverberg et al., 1982; Song et al., 1996; Lee et al., 1999c; Vais et al., 2000). Cohen and colleagues (Vais et al., 2000) compared two tail current recording protocols, one with a 500 ms depolarization from –100 to 0 mV and the other with a 100-pulse train of 5 ms depolarization from –100 to 0 mV. They found that the train of pulses produced a more pronounced tail current, suggesting that deltamethrin interacts with the open state of the channel (Vais et al., 2000). Therefore, for measurement of pyrethroid-induced tail current in this




Fig. 1. Functional expression of a cockroach  $\text{Para}^{\text{CSMA}}$  sodium channel in *Xenopus* oocytes (coexpressed with *D. melanogaster tipE*). (A) Sodium current traces. Sodium currents were recorded from a holding potential of  $-120$  mV during depolarizations from  $-80$  to  $65$  mV in  $5$  mV increments. (B) Dose-dependent block of sodium current by TTX. Amplitudes of the peak currents measured in the presence of various concentrations of TTX were normalized using the amplitude of the maximal peak current in the absence of TTX. The data were fitted with a sigmoid curve;  $\text{EC}_{50}$  is  $0.9$  nM. Symbols represent means and error bars indicate the SEM for five oocytes. (C) Sodium current–voltage relation ( $I$ – $V$  curve). Current were recorded as described for (A), and the peak current amplitude is plotted versus the depolarization potential. Symbols represent means and error bars indicate the SEM for five oocytes. (D) Normalized conductance–voltage ( $G$ – $V$ ) and inactivation curves. The peak current in (C) was converted to conductance as described in Section 2 and plotted against the depolarizing voltage (triangles). The voltage-dependence of inactivation was determined using  $200$  ms inactivating pre-pulses from a holding potential of  $-120$  to  $+40$  mV in  $5$  mV increments, followed by test pulses to  $-5$  mV for  $12$  ms. The peak current amplitude during the test depolarization was normalized to the maximum current amplitude, and plotted as a function of the pre-pulse potential (circles). The smooth curves represent the best fit using Boltzmann equations, as described in Section 2. Symbols represent means; error bars indicate the SEM from three oocytes.

study, we adopted the recording protocol of a 100-pulse train of  $5$  ms depolarization from  $-120$  to  $0$  mV with a  $2$ -ms interval between each depolarization. Traces of tail current were recorded  $5$  min after application of each deltamethrin concentration. The deltamethrin-induced tail current increased initially and then gradually decreased, generating a hooked tail current (Fig. 3).

The minimum effective deltamethrin concentration required to elicit a detectable tail current is approximately  $0.01$   $\mu\text{M}$  for the  $\text{Para}^{\text{CSMA}}$ ,  $\text{Para}^{\text{CSMA-K}}$  and  $\text{Para}^{\text{CSMA-R}}$  channels (Fig. 3A–C). A large tail current was elicited at  $1.0$   $\mu\text{M}$  (Fig. 3A–C). Concentrations above  $1.0$   $\mu\text{M}$  resulted in large leakage currents that made it impossible to clamp the oocytes. Conversely, much higher concentrations of deltamethrin ( $10$ –

$100$   $\mu\text{M}$ ) were required to elicit significant tail currents for  $\text{Para}^{\text{CSMA-F}}$ ,  $\text{Para}^{\text{CSMA-KF}}$ ,  $\text{Para}^{\text{CSMA-RF}}$  and particularly  $\text{Para}^{\text{CSMA-KRF}}$  channels (Fig. 3D, F–H), clearly demonstrating the importance of these mutations in  $\text{Para}^{\text{CSMA}}$  channel insensitivity to deltamethrin in the German cockroach. Surprisingly the  $\text{Para}^{\text{CSMA-KR}}$  channel, which carries the two novel mutations E434K and C764R but not the L993F mutation, was more sensitive to deltamethrin than the wild-type channel or channels carrying only the E434K or C764R single mutation (Fig. 3E).

To quantify the differences in sensitivity to deltamethrin among channels, we calculated the percentage of channels modified by deltamethrin as described in Section 2 (Tatebayashi and Narahashi, 1994). The percentage modification was plotted as a function of the

Table 1  
Voltage dependence of activation and inactivation of Para<sup>CSMA</sup> sodium channels

Na <sup>+</sup> channel type	Mutations	Activation <sup>a</sup>		Inactivation <sup>a</sup>	
		V <sub>1/2</sub> (mV)	k (mV)	V <sub>1/2</sub> (mV)	k (mV)
Para <sup>CSMA</sup>	Wild-type	-29.0±1.3	5.5±0.5	-42.4±0.4	5.0±0.1
Para <sup>CSMA-F</sup>	L993F	-27.8±2.4	6.2±0.4	-44.0±1.9	5.3±0.4
Para <sup>CSMA-R</sup>	C764R	-27.9±4.5	5.6±0.6	-43.8±1.2	5.6±0.3
Para <sup>CSMA-K</sup>	E434K	-25.7±3.7	5.4±1.0	-43.2±0.9	5.4±0.3
Para <sup>CSMA-KF</sup>	E434K, L993F	-30.3±4.1	5.1±0.7	-39.3±1.1	5.1±0.8
Para <sup>CSMA-RF</sup>	C764R, L993F	-24.2±2.7	5.3±0.7	-40.6±0.4	5.1±0.3
Para <sup>CSMA-KR</sup>	E434K, C764R	-24.5±4.8	5.7±0.6	-44.4±3.2	5.6±0.2
Para <sup>CSMA-KRF</sup>	E434K, C764R, L993F	-27.9±8.3	5.9±0.8	-40.9±1.7	5.2±0.8

<sup>a</sup> The voltage-dependence of conductance and inactivation data were fit with two-state Boltzmann equations, as described in Section 2, to determine V<sub>1/2</sub>, the voltage for half-maximal conductance or inactivation and k, the slope factor for conductance or inactivation. Each value represents the mean ± SEM for three oocytes.




Fig. 2. Schematic drawing of the cockroach Para<sup>CSMA</sup> sodium channel protein indicating kdr-associated mutations. Four homologous domains (I–IV) and six transmembrane segments (S1–6) are shown. The three kdr-associated mutations (E434K, C764R, L993F) used in this study are indicated.

Table 2  
Peak current amplitudes of wild-type and mutant Para<sup>CSMA</sup> sodium channels in *Xenopus* oocytes

Na <sup>+</sup> channel type	I <sub>max</sub> (μA) <sup>a</sup>	N <sup>b</sup>
Para <sup>CSMA</sup>	5.59±1.20	12
Para <sup>CSMA-F</sup>	3.89±1.44	6
Para <sup>CSMA-R</sup>	0.96±0.24	4
Para <sup>CSMA-K</sup>	1.29±0.23	5
Para <sup>CSMA-RF</sup>	5.77±1.02	14
Para <sup>CSMA-KR</sup>	5.90±1.17	9
Para <sup>CSMA-KF</sup>	5.98±0.83	15
Para <sup>CSMA-KRF</sup>	7.04±1.26	6

<sup>a</sup> Amplitude of the maximal peak current measured during a 20 ms depolarization from -120 to -10 mV.

<sup>b</sup> Number of oocytes.

deltamethrin concentration (Fig. 4). Modification of 20% of sodium channels required five-fold more deltamethrin for the Para<sup>CSMA-F</sup> channel, 100-fold more for the Para<sup>CSMA-KF</sup> and Para<sup>CSMA-RF</sup> channels, and 500-fold more for the Para<sup>CSMA-KRF</sup> channel, compared to the wild-type channel. The sensitivity of the Para<sup>CSMA-R</sup> channel to deltamethrin was similar to that of the wild-

type channel. Para<sup>CSMA-K</sup> and Para<sup>CSMA-KR</sup> channels were two-fold and 10-fold, respectively, more sensitive to deltamethrin than the wild-type channel (Fig. 4).

The tail currents of Para<sup>CSMA-KF</sup>, Para<sup>CSMA-RF</sup> and Para<sup>CSMA-KRF</sup> channels clearly decayed more completely than the tail currents of the other channels including the Para<sup>CSMA-F</sup> channel (Fig. 3). To quantify this difference, the largest tail current in Fig. 3 was used for the analysis of tail current decay for each channel. The tail current decay was best fitted with a single exponential equation. The time constants for Para<sup>CSMA-KF</sup>, Para<sup>CSMA-RF</sup> and Para<sup>CSMA-KRF</sup> channels were significantly smaller than those of the other channels (Table 3).

#### 4. Discussion

This is the first functional characterization of a cockroach sodium channel Para<sup>CSMA</sup> expressed in *Xenopus* oocytes. Similar to the *D. melanogaster* Para and house fly Vssc1 sodium channels, the cockroach Para<sup>CSMA</sup> sodium channel is TTX-sensitive (an EC<sub>50</sub> of 0.9 nM) and requires *Drosophila tipE*, a potential subunit of




Fig. 3. *Para<sup>CSMA</sup>* channel sensitivity to deltamethrin. (A)–(H) Tail currents induced by deltamethrin in oocytes expressing the wild-type and mutant *Para<sup>CSMA</sup>* channels. Tail currents were recorded in response to a 100-pulse train of 5 ms depolarizations from  $-120$  to  $0$  mV with a 2-ms interval between each depolarization. K, R, F, KR, KF, RF, KRF represent sodium channels carrying E434K, C764R, L993F, E434K+C764R, E434K+L993F, C764R+L993F and E434K+C764R+L993F, respectively. Note the differences in the scales of tail current and also the concentrations of deltamethrin. At  $100 \mu$ M, the recording solution showed a slight cloudiness, indicating that deltamethrin was not completely soluble at this concentration.




Fig. 4. Quantification of Para<sup>CSMA</sup> and mutant channel sensitivity to deltamethrin. The percentage of channels modified by deltamethrin was determined using the equation  $M = [I_{tail}/(E_h - E_{Na})]/[I_{Na}/(E_t - E_{Na})] \times 100$  (Tatebayashi and Narahashi, 1994) (see Section 2). Percentage of modification of sodium channels is plotted as a function of deltamethrin concentration. The data were fitted with the Hill equation (see Section 2). EC<sub>20</sub> values (inset) are derived from the fitted curves. Hill coefficient ranges 0.8–1.3 for different channels, except for the KRF channel, which gives a Hill coefficient of 0.4. Each point represents mean  $\pm$  SEM (pooled data from five oocytes).

Table 3  
Time constant of tail current decay induced by deltamethrin

Na <sup>+</sup> channel type	$\tau^a$ (ms)	N <sup>b</sup>
Para <sup>CSMA</sup>	1537 $\pm$ 737	20
Para <sup>CSMA</sup> -F	1408 $\pm$ 353	12
Para <sup>CSMA</sup> -R	1518 $\pm$ 227	7
Para <sup>CSMA</sup> -K	1679 $\pm$ 328	7
Para <sup>CSMA</sup> -RF	1102 $\pm$ 77 <sup>c</sup>	11
Para <sup>CSMA</sup> -KR	1677 $\pm$ 357	14
Para <sup>CSMA</sup> -KF	1034 $\pm$ 96 <sup>c</sup>	7
Para <sup>CSMA</sup> -KRF	1042 $\pm$ 130 <sup>c</sup>	9

<sup>a</sup> The largest deltamethrin-induced tail currents (see Fig. 3) was fitted with a single exponential equation, as described in Section 2. Each value represents the mean  $\pm$  SEM.

<sup>b</sup> Number of oocytes.

<sup>c</sup> Indicates a statistically significant difference compared to the wild-type channel at  $p < 0.05$ .

insect sodium channels (Feng et al., 1995; Warmke et al., 1997), for robust expression. Most importantly, we demonstrated a critical role of two novel *para*<sup>CSMA</sup> gene mutations (E434K and C764R) in reduced Para<sup>CSMA</sup> channel sensitivity to deltamethrin. Specifically, we found that: (i) the previously characterized L993F mutation alone conferred only a low level of insensitivity to deltamethrin; (ii) E434K or C764R alone significantly reduced sodium peak currents but had no effect on Para<sup>CSMA</sup> channel sensitivity to deltamethrin; and (iii) when E434K and C764R were combined with L993F (i.e. Para<sup>CSMA</sup>-KF, Para<sup>CSMA</sup>-RF and Para<sup>CSMA</sup>-KRF channels), however, sodium current magnitudes were

restored to the wild-type level and channel sensitivity to deltamethrin was markedly reduced. These results confirm our previous speculation (Liu et al., 2000) that the E434K and C764R mutations, in the presence of L993F, are responsible for high levels of pyrethroid-resistance in the German cockroach.

In the house fly, the super-kdr mutation (M918T) in the linker connecting IIS4 and IIS5, when combined with the L to F mutation (L1014F in house fly), confers higher-level resistance to pyrethroids (Lee et al., 1999c). This super-kdr mutation is also present in pyrethroid-resistant horn flies (Guerrero et al., 1997). The fact that highly pyrethroid-resistant cockroaches contain the E434K or C764R mutations instead of the M918T mutation is somewhat unexpected. This variation in the mutations that cause higher levels of resistance is in contrast to the absolute conservation of the kdr-associated mutation (L1014F in house fly, and L993F in German cockroach in IIS6) in all characterized kdr and kdr-type insects (Williamson et al., 1996; Miyazaki et al., 1996; Dong, 1997; Guerrero et al., 1997; Martinez-Torres et al. 1997, 1998; Martinez-Torres et al., 1999a,b; Lee et al., 1999a). Therefore, these results establish that different sodium channel mutations, when combined with the L1014F mutation in house fly or the L993F mutation in German cockroach, can result in high-level resistance to pyrethroids. It remains an interesting question why distinct secondary mutations are selected in house fly and cockroach to enhance the level of pyrethroid resistance.

It is important to note that although L993F alone reduced channel sensitivity to a pyrethroid, as was observed for the house fly kdr mutation (L1014F) and the *Heliothis virescens* L1029H mutation, neither the E434K mutation nor the C764R mutation alone reduced Para<sup>CSMA</sup> channel sensitivity to pyrethroids. This is in sharp contrast to the house fly M918T mutation, which alone reduces sodium channel sensitivity to pyrethroids (Lee et al., 1999c). In fact, the Para<sup>CSMA</sup> channel bearing the E434K and C764R double mutation is more sensitive to deltamethrin (Fig. 3). Furthermore, we did not detect significant alteration of channel gating properties by any of the three single mutations or various mutation combinations. The housefly L1014F mutation, however, causes a depolarizing shift (ca. 6-mV) of voltage-dependence of activation (Lee et al., 1999c). A similar shift was observed when the *Heliothis virescens* L1029H mutation was introduced into the *Drosophila* Para channel (Zhao et al., 2000). However, when the housefly L1014F mutation was introduced into the *Drosophila* Para channel, no significant shift was observed (Vais et al., 2000). Our results are consistent with the latter.

The hooked tail currents induced by deltamethrin in this study have been observed in mammalian neurons and sodium channels expressed in oocytes (such as Lund and Narahashi, 1982; Vijverberg et al., 1982; Song et al., 1996; Vais et al., 2000). The initial increasing phase

of the tail current is the result of the delayed closing of the inactivation gate upon termination of depolarization. The subsequent decay of the tail current reflects the slow rate of deactivation of the modified channels. The mutation combinations, RF, KF and KRF, not only reduce the channel sensitivity to deltamethrin, also cause more complete decay of the tail current possibly by increasing the rate of deactivation of the modified channels. An acceleration of tail current decay was previously observed for the house fly *kdr* and *super-kdr* mutations and the *Heliothis virescens* L1029H mutation (Smith et al., 1997; Vais et al., 2000; Zhao et al., 2000). However, the cockroach L993F mutation reduces the channel sensitivity to deltamethrin without acceleration of the decay of the tail current (Table 3 and Fig. 3).

An intriguing observation from this study is that the E434K or C764R mutation alone significantly reduced the peak sodium current by more than four-fold, whereas the L993F mutation alone did not affect the peak sodium current or channel gating (Tables 1 and 3). Reduction of peak sodium currents was also observed for house fly sodium channels carrying the M918T mutation (Lee et al., 1999c). The neutral effects of the L993F mutation alone on sodium channel peak current and gating can explain the genetic stability and presence of this mutation in all characterized *kdr* and *kdr*-type insect pest species/populations collected from diverse regions. However, the negative effect of the E434K or C764R mutation on peak sodium current indicates that insects carrying either of these two mutations may have decreased fitness. In support of this speculation, we have not detected any cockroaches containing only the E434K or C764R mutation (Liu et al., 2000; Z. Liu and K. Dong, unpublished data). The mechanism by which the E434K and C764R mutations decrease sodium current amplitudes is not known. It is possible that these substitutions alter the folding or assembly of the Para<sup>CSMA</sup> protein, which would result in fewer functional channels, or they might decrease either the single channel conductance or probability of channel opening, or they decrease the interaction of Para<sup>CSMA</sup> with TipE.

Interestingly, when E434K or C764R is combined with L993F, normal peak current amplitudes were observed (Table 3). Similarly, the concomitant expression of the L1014F and M918T mutations in the house fly sodium channel results in normal peak current amplitudes (Lee et al., 1999c). Based on this observation, we speculate that an initial selection of the L993F mutation (most likely by DDT; see Dong, 1997) must have provided a platform for subsequent selection of the E434K and C764R mutations by pyrethroids in the German cockroach. Presence of the L993F mutation in the *para*<sup>CSMA</sup> gene would compensate for the functional constraint imposed by E434K and C764R mutations. Therefore, the L993F mutation must have played a crucial role in the evolution of high-level pyrethroid resistance in

cockroaches, allowing selection of additional *para*<sup>CSMA</sup> gene mutations that otherwise would decrease the function of the Para<sup>CSMA</sup> channel. Furthermore, we found that the Para<sup>CSMA-KR</sup> channel (double mutations in the absence of L993F) was more susceptible to deltamethrin (Fig. 3) and would unlikely have survived pyrethroid selection in the field. All these observations suggest a 'successive mutation' mechanism by which German cockroaches adapted to pyrethroid selective pressure. A similar mechanism has been proposed by Lee et al. (1999c) to explain the selection of the *super-kdr* mutation (M918T) in the housefly. Such a mechanism is in contrast to the 'mutation recombination' mechanism, in which preexisting intragenic resistance mutations in the acetylcholinesterase gene are recombined to produce higher levels of resistance to organophosphates (Mutero et al., 1994). Thus, the successive mutation mechanism appears to represent an alternative paradigm for evolution of insecticide resistance.

## Acknowledgements

We thank R. Hollingworth, N. Koller and anonymous reviewers for critical review of the manuscript, and C.J. Cohen for helpful discussions during the course of this work and sharing unpublished results. This work was supported in part by National Science Foundation grants, IBN 9696092 and IBN 9808156 (to K.D.), National Institutes of Health Grant 08-1GM57440A (to K.D.) and the Michigan State University Rackham Endowment Fund (to K.D.).

## References

- Bloomquist, J.R., Soderlund, D.M., 1988. Pyrethroid insecticides and DDT modify alkaloid-dependent sodium channel activation and its enhancement by sea anemone toxin. *Mol. Pharmacol.* 33, 543–550.
- Brown, G.B., Gaupp, J.E., Olsen, R.W., 1988. Pyrethroid insecticides: Stereospecific allosteric interaction with the [<sup>3</sup>H]-batrachotoxinin A-20-a-benzoate binding site of mammalian voltage-sensitive sodium channels. *Mol. Pharmacol.* 34, 54–59.
- Catterall, W.A., 1988. Molecular pharmacology of voltage-sensitive sodium channels. *ISI Atlas Sci. Pharmacol.* 190–195.
- Catterall, W.A., 1992. Cellular and molecular biology of voltage-gated sodium channels. *Physiol. Rev.* 72, S15–S48.
- Dong, K., 1997. A single amino acid change in the Para sodium channel protein is associated with knockdown-resistance (*kdr*) to pyrethroid insecticides in German cockroach. *Insect Biochem. Mol. Biol.* 27, 93–100.
- Dong, K., Valles, S.M., Scharf, M.E., Zeichner, B., Bennett, G.W., 1998. Knockdown resistance (*kdr*) mutation in pyrethroid-resistant German cockroaches. *Pestic. Biochem. Physiol.* 60, 195–204.
- Feng, G., Deak, P., Chopra, M., Hall, L.M., 1995. Cloning and functional analysis of TipE, a novel membrane protein that enhances *Drosophila* Para sodium channel function. *Cell* 82, 1001–1011.
- Goldin, A.L., 1992. Maintenance of *Xenopus laevis* and oocyte injection. *Methods Enzymol.* 207, 266–279.
- Guerrero, F.D., Jamroz, R.C., Kammlah, D., Kunz, S.E., 1997. Toxicology

- logical and molecular characterization of pyrethroid-resistant horn flies, *Haematobia irritans*: Identification of kdr and super-kdr point mutations. *Insect Biochem. Mol. Biol.* 27, 745–755.
- Hille, B., 1992. Modifiers of gating. In: Hille, B. (Ed.), *Ionic Channels of Excitable Membranes*. Sinauer Associates, Massachusetts, pp. 472–503.
- Jacques, Y., Romey, G., Carvey, M.T., Kartalovski, B., Lazdunski, M., 1980. Interaction of pyrethroids with the Na<sup>+</sup> channel in mammalian neuronal cells in culture. *Biochim. Biophys. Acta* 600, 882–897.
- Kontis, K.J., Goldin, A.L., 1993. Site-directed mutagenesis of the putative pore region of the rat IIA sodium channel. *Mol. Pharmacol.* 43, 635–644.
- Lee, S.H., Dunn, J.B., Clark, J.M., Soderlund, D.M., 1999a. Molecular analysis of kdr-like resistance in a permethrin-resistant strain of Colorado potato beetle. *Pestic. Biochem. Physiol.* 63, 63–75.
- Lee, D., Park, Y., Brown, T.M., Adams, M.E., 1999b. Altered properties of neuronal sodium channels associated with genetic resistance to pyrethroids. *Mol. Pharmacol.* 55, 584–593.
- Lee, S.H., Smith, T.J., Knipple, D.C., Soderlund, D.M., 1999c. Mutations in the house fly *Vssc1* sodium channel gene associated with super-kdr resistance abolish the pyrethroid sensitivity of *Vssc1*/tipE sodium channels expressed in *Xenopus* oocytes. *Insect Biochem. Mol. Biol.* 29, 185–194.
- Liu, Z., Valles, S.M., Dong, K., 2000. Novel point mutations in the German cockroach *para* sodium channel gene are associated with knockdown resistance (kdr) to pyrethroid insecticides. *Insect Biochem. Mol. Biol.* 30, 991–997.
- Lombet, A., Mourre, C., Lazdunski, M., 1988. Interaction of insecticides of the pyrethroid family with specific binding sites on the voltage-dependent sodium channel from mammalian brain. *Brain Res.* 459, 44–53.
- Lund, A.E., Narahashi, T., 1982. Dose-dependent interaction of the pyrethroid isomers with sodium channels of squid axon membranes. *NeuroTox.* 3, 11–24.
- Martinez-Torres, D., Devonshire, A.L., Williamson, M.S., 1997. Molecular studies of knockdown resistance to pyrethroids: cloning of domain II sodium channel sequences from insects. *Pestic. Sci.* 51, 265–270.
- Martinez-Torres, D., Chandre, F., Williamson, M.S., Darriet, F., Berge, J.B., Devonshire, A.L., Guillet, P., Pasteur, N., Pauron, D., 1998. Molecular characterization of pyrethroid knockdown resistance (kdr) in the major malaria vector *Anopheles gambiae*. *Insect Mol. Biol.* 7, 179–184.
- Martinez-Torres, D., Chevillon, C., Brun-Barale, A., Berge, J.B., Pasteur, N., Pauron, D., 1999a. Voltage-dependent Na<sup>+</sup> channels in pyrethroid-resistant *Culex pipiens* L. mosquitoes. *Pestic. Sci.* 55, 1012–1020.
- Martinez-Torres, D., Foster, S.P., Field, L.M., Devonshire, A.L., Williamson, M.S., 1999b. A sodium channel point mutation is associated with resistance to DDT and pyrethroid insecticides in the peach-potato aphid, *myzus persicae*. *Insect Mol. Biol.* 8, 339–346.
- Miyazaki, M., Ohyama, K., Dunlap, D.Y., Matsumura, F., 1996. Cloning and sequencing of the *para*-type sodium channel gene from susceptible and kdr-resistant German cockroaches (*Blattella germanica*) and house fly (*Musca domestica*). *Mol. Gen. Genet.* 252, 61–68.
- Mutero, A., Pralavorio, M., Bride, J.M., Fournier, D., 1994. Resistance-associated point mutations in insecticide-insensitive acetylcholinesterase. *Proc. Natl. Acad. Sci. US A* 91, 5922–5926.
- Narahashi, T., 1986. Toxins that modulate the sodium channel gating mechanism. In: Kao, C.Y., Levinsonin, S.R. (Eds.), *Tetrodotoxin, Saxitoxin and the Molecular Biology of the Sodium Channel*. Annals New York Academy of Sciences, New York, pp. 133–151.
- Narahashi, T., 1988. Molecular and cellular approaches to neurotoxicology: past, present and future. In: Lunt, G.G. (Ed.), *Neurotox '88: Molecular Basis of Drug and Pesticide Action*. Elsevier, New York, pp. 563–582.
- Park, Y., Taylor, M.F., 1997. A novel mutation L1029H in sodium channel *hscp* associated with pyrethroid resistance for *Heliothis virescens* (Lepidoptera: Noctuidae). *Insect Biochem. Mol. Biol.* 27, 9–13.
- Park, Y., Taylor, M.F., Feyereisen, R., 1997. A valine421 to methionine mutation in IS6 of the *hscp* voltage-gated sodium channel associated with pyrethroid resistance in *Heliothis virescens*. *Biochem. Biophys. Res. Commun.* 239, 688–691.
- Smith, T.J., Lee, S.H., Ingles, P.J., Knipple, D.C., Soderlund, D.M., 1997. The L1014F point mutation in the House Fly *Vssc1* sodium channel confers knockdown resistance to pyrethroids. *Insect Biochem. Mol. Biol.* 27, 807–812.
- Soderlund, D.M., Bloomquist, J.R., 1989. Neurotoxic actions of pyrethroid insecticides. *Annu. Rev. Entomol.* 34, 77–96.
- Soderlund, D.M., Bloomquist, J.R., 1990. Molecular mechanisms of insecticide resistance. In: Roush, R.T., Tabashnikin, B.E. (Eds.), *Pesticide Resistance in Arthropods*. Chapman and Hall, New York, pp. 58–96.
- Song, J.H., Nagata, K., Tatebayashi, H., Narahashi, T., 1996. Interactions of tetramethrin, fenvalerate and DDT at the sodium channel in rat dorsal root ganglion neurons. *Brain Res.* 708, 29–37.
- Takeda, K., Narahashi, T., 1988. Chemical modification of sodium channel inactivation: separate sites for the action of grayanotoxin and tetramethrin. *Brain Res.* 448, 308–312.
- Tatebayashi, H., Narahashi, T., 1994. Differential mechanism of action of the pyrethroid tetramethrin on tetrodotoxin-sensitive and tetrodotoxin-resistant sodium channels. *J. Pharmacol. Exp. Ther.* 270, 595–603.
- Trainer, V.L., McPhee, J.C., Boutelet-Bochan, H., Baker, C., Scheuer, T., Babin, D., Demoute, J.P., Guedin, D., Catterall, W.A., 1997. High affinity binding of pyrethroids to the  $\alpha$ -subunit of brain sodium channel. *Mol. Pharmacol.* 51, 651–657.
- Vais, H., Williamson, M.S., Goodson, S.J., Devonshire, A.L., Warmke, J.W., Usherwood, P.N.R., Cohen, C., 2000. Activation of *Drosophila* sodium channels promotes modification by deltamethrin: reductions in affinity caused by knock-down resistance mutations. *J. Gen. Physiol.* 115, 305–318.
- Vijverberg, H.P.M., van der Zalm, J.M., van den Bercken, J., 1982. Similar mode of action of pyrethroids and DDT on sodium channel gating in myelinated nerves. *Nature* 295, 601–603.
- Warmke, J.W., Reenan, R.A.G., Wang, P., Qian, S., Arena, J.P., Wang, J., Wunderler, D., Liu, K., Kaczorowski, G.J., Van Der Ploeg, L.H.T., Ganetzky, B., Cohen, C.J., 1997. Functional expression of *Drosophila para* sodium channels: Modulation by the membrane protein tipE and toxin Pharmacology. *J. Gen. Physiol.* 110, 119–133.
- Williamson, M.S., Martinez-Torres, D., Hick, C.A., Devonshire, A.L., 1996. Identification of mutations in the housefly *para*-type sodium channel gene associated with knockdown resistance (kdr) to pyrethroid insecticides. *Mol. Gen. Genet.* 252, 51–60.
- Zhao, Y., Park, Y., Adams, M.E., 2000. Functional and evolutionary consequences of pyrethroid resistance mutations in S6 transmembrane segments of a voltage-gated sodium channel. *Biochem. Biophys. Res. Commun.* 278, 516–521.
- Zlotkin, E., 1999. The insect voltage-gated sodium channel as target of insecticides. *Annu. Rev. Entomol.* 44, 429–455.